


# Gouget N

Wine grape variety.



## Origin

This variety was originally from the center of France, seemingly from the Montluçon region. Based on published genetic analyses, Gouget is probably a descendant of Gouais blanc.

## Use

Wine grape variety.

## Name of the variety in France

Gouget

## Synonymy

There is no officially recognized synonym in France nor in the other countries of the European Union, for this variety.

## Regulatory data

In France, Gouget is officially listed in the "Catalogue of vine varieties" on the A list and classified.

## Description elements

The identification is based on:

- the tip of the young shoot with a low density of prostrate hairs,
- the green young leaves with bronze spots,
- the adult leaves, entire or with three lobes, with an open V-shaped petiole sinus, no anthocyanin coloration of veins, an involute leaf blade, and on the lower side of the leaves, a medium density of erect hairs,
- the round-shaped berries.

# Evolution of cultivated areas in France

Year ha

1958739

1968518

197913

19884

199816

200813

20188.4

## Genetic profile

Microsatellite	VVS2	VVMD5	VVMD7	VVMD27	VRZAG62	VRZAG79	VVMD25	VVMD28	VVMD32
Allele 1	131	223	239	176	194	238	238	227	251
Allele 2	149	232	257	191	196	252	248	233	271

### Cultivation and agronomic skills

This variety is vigorous with an erect bearing and is generally pruned long. Fruit production is good after spring frost.

### Susceptibility to Diseases and Pests

Gouget is very susceptible to grey rot.

### Clonal selection in France

There is no certified clone for this variety yet.

### Bibliographic references

- Catalogue des variétés et clones de vigne cultivés en France. Collectif, 2007, Ed. IFV, Le Grau-du-Roi, France.
- Documentary collections of the Centre de Ressources Biologiques de la Vigne de Vassal-Montpellier, INRAE - Institut Agro | Montpellier, Marseillan, France.
- Dictionnaire encyclopédique des cépages et de leurs synonymes. P. Galet, 2015, Ed. Libre&Solidaire, France.
- Traité général de viticulture, Ampélographie. P. Viala and V. Vermorel, 1901-1909, Ed. Masson, Paris, France.

### Phenology

Bud burst: 2 days after Chasselas.

Grape maturity: early-season, 1 week after Chasselas.

### Technological potential

The bunches and berries are small. This variety produces relatively acidic, not very colored and not very tannic wine, with low alcohol level.



---

*Plantgrape, all rights reserved,  
plantgrape.com, © UMT Géno-Vigne®  
INRA - IFV - L'Institut Agro Montpellier*