

Mourvèdre N

Wine grape variety.



Origin

This variety is originally from Spain and was probably introduced in France at the end of the Middle Ages.

Use

Wine grape variety.

Name of the variety in France

Mourvèdre

Synonymy

In the European Union, Mourvèdre is officially called by other names: Mataro (Cyprus) and Monastrell (Spain). These synonyms are officially recognized in France regarding plant propagation material.

Regulatory data

In France, Mourvèdre is officially listed in the "Catalogue of vine varieties" on the A list and classified. This variety is also listed in the catalogues of other Member States of the European Union: Croatia, Cyprus, Greece, Malta and Spain.

Description elements

The identification is based on:

- the tip of the young shoot with a very high density of prostrate hairs,
- the young leaves with bronze spots,
- the shoots with red colored internodes and an erect bearing,
- the circular adult leaves, entire or with three lobes, with a open lyre-shaped petiole sinus, large teeth, moderate compared to their width at the base with straight sides, a moderate anthocyanin coloration of veins, a flat leaf blade, and on the lower side of the leaves, a high density of erect and prostrate hairs,
- the round-shaped berries.

Evolution of cultivated areas in France

Year ha

1958617

1968857

19793146

19885608

19987249

20089164

20189149

Genetic profile

| Microsatellite | VVS2 | VVMD5 | VVMD7 | VVMD27 | VRZAG62 | VRZAG79 | VVMD25 | VVMD28 | VVMD32 |
|----------------|------|-------|-------|--------|---------|---------|--------|--------|--------|
| Allele 1 | 131 | 223 | 249 | 176 | 188 | 252 | 240 | 243 | 239 |
| Allele 2 | 149 | 238 | 249 | 186 | 204 | 262 | 262 | 257 | 255 |

Cultivation and agronomic skills

Mourvèdre is a demanding variety and its growing is delicate. With its erect bearing and acrotonic nature, it must be pruned short. As such, it can be trained in cordon, with reduced trellis or in open gobelet. Care must be taken to maintain the non abundant foliage, that is moderately effective. Trimming must be avoided to limit the presence of large size grappillons, which then represent a bothersome competition. Production per vine trunk must be low to prevent risk of depletion. This variety sometimes demonstrates alternating fertility. The pedo-climatic requirements of Mourvèdre are very demanding. This variety appears well adapted to limestone soils deep enough to allow a limited but regular water supply. Potassium and magnesium supplies (K/Mg) must also be balanced in order to prevent the appearance of deficiencies. Finally, the main limiting factor for growing this late variety (budburst and grape maturity), is that the temperatures at the moment of grape maturation must be very high (especially minimum temperatures) to enable proper ripening and a sufficient accumulation of sugar in the berries. This variety is thus reserved for the hottest southern areas and the best exposed.

Susceptibility to Diseases and Pests

Mourvèdre is sensitive to mites, vine leafhoppers, esca and sour rot. On the other hand, it is generally not very so susceptible to grey rot and it resists well against phomopsis.

Clonal selection in France

The fourteen certified Mourvèdre clones carry the numbers 233, 234, 244, 245, 247, 248, 249, 369, 448, 449, 450, 520, 1069 and 1215. A conservatory of almost 100 clones has been planted in the wine-growing region of Bandol (French department of Var) since 2004.

Bibliographic references

- Catalogue des variétés et clones de vigne cultivés en France. Collectif, 2007, Ed. IFV, Le Grau-du-Roi, France.
- Documentary collections of the Centre de Ressources Biologiques de la Vigne de Vassal-Montpellier, INRAE - Institut Agro | Montpellier, Marseillan, France.
- Dictionnaire encyclopédique des cépages et de leurs synonymes. P. Galet, 2015, Ed. Libre&Solidaire, France.
- Traité général de viticulture, Ampélographie. P. Viala and V. Vermorel, 1901-1909, Ed. Masson, Paris, France.

Phenology

Bud burst: 13 days after Chasselas.

Grape maturity: late-season, 4 weeks and a half after Chasselas.

Technological potential

Mourvèdre's bunches are medium to large and the berries are medium in size. The color potential of this variety is moderate with sometimes a lack of acidity. Nevertheless, Mourvèdre can produce very high quality, aromatic, tannic and structured wines, suitable for ageing and maturing in wood barrels, provided that the grape maturity is sufficient (12 natural degrees minimum) and that the production per vine trunk is limited.



