

Raisaine B

Wine grape variety.



Origin

This variety is originally from the Cévennes ardéchoises. Based on genetic analysis carried out in Montpellier, it would result from the crossbreeding of Ribier noir and Grec rouge.

Use

Wine grape variety.

Name of the variety in France

Raisaine

Synonymy

There is no officially recognized synonym in France nor in the other countries of the European Union, for this variety.

Regulatory data

In France, Raisaine is officially listed in the "Catalogue of vine varieties" since 2016 on the A and classified.

Description elements

The identification is based on:

- the tip of the young shoot with a medium density of prostrate hairs and a piping distribution of anthocyanin coloration,
- the green yellow young leaves with bronze spots and a low density of prostrate hairs,
- the shoots with short, red or red-striped internodes,
- the wedge-shaped or circular, dark green adult leaves, with five lobes, deep V-shaped upper lateral sinuses, an open or slightly open V- or brace-shaped petiole sinus, moderately long teeth compared to their width at the base with straight sides or with one side concave and one side convex, a weak anthocyanin coloration of veins, a thick, finely blistered, twisted, sometimes involute leaf blade, and on the lower side of the leaves, no or a very low density of erect and prostrate hairs,
- the round-shaped berries with rather numerous brown lenticels on the surface.

Evolution of mother vine surfaces

Year

2018

ha

0.1

Genetic profile

Microsatellite	VVS2	VVMD5	VVMD7	VVMD27	VRZAG62	VRZAG79	VVMD25	VVMD28	VVMD32
Allele 1	131	223	247	178	194	244	240	245	261
Allele 2	131	234	249	182	204	252	240	257	271

Cultivation and agronomic skills

Raisaine is vigorous, productive and has a semi-erect or erect bearing. It can be pruned short.

Clonal selection in France

The only certified Raisaine clone carries the number 1297.

Phenology

Bud burst: 4 days before Chasselas.

Grape maturity: early-season, 1 week and a half to 2 weeks after Chasselas.


Technological potential

Raisaine's bunches are medium in size and compact. The berries are medium to large, simple-flavored, with a moderately thick skin, a lot of bloom and a firm and juicy pulp. Raisaine produces round, fluid wines with a low acidity and honey and quince aromas. Raisaine could also be used as table grapes.


Susceptibility to Diseases and Pests

Raisaine is sensitive to downy mildew and less to powdery mildew.

Debourrement



Maturité



Bibliographic references

- Documentary collections of the Centre de Ressources Biologiques de la Vigne de Vassal-Montpellier, INRAE -
©2023 Plantgrape, all rights reserved

Institut Agro Montpellier, Marseillan, France.

- Traité général de viticulture, Ampélographie. P. Viala and V. Vermorel, 1901-1909, Ed. Masson, Paris, France.

- Le vignoble. A. Mas and V. Pulliat, 1874-1879, Ed. Masson, Paris, France.

- Registre Ampélographique International. 1961-1972, Ed. OIV, France.



*Plantgrape, all rights reserved,
plantgrape.fr, UMT Géno-Vigne®
INRAE - IFV - L'Institut Agro Montpellier*