

# New

# Patricia INTA Rs

Grape variety intended for the production of fruit juices.



## Origin

This variety has been obtained by the INTA Station of Mendoza (Argentina) by A. Gargiulo by crossbreeding Moscatel rosado by a descendant of Gibi and Sultanine.

#### Use

Grape variety intended for the production of fruit juices.

Name of the variety in France

Patricia INTA

#### Synonymy

There is no officially recognized synonym in France nor in the other countries of the European Union, for this variety.

#### **Regulatory data**

In France, Serna INTA is officially listed in the "Catalogue of vine varieties" on the A list since 2020.

#### **Description elements**

The identification is based on:

- the tip of the young shoot with a low density of prostrate hairs,
- the green young leaves with a low density of prostrate hairs,
- the green shoots,

- the rather wide adult leaves, with five leaves, shallow lower lateral sinuses, an open U- or V-shaped petiole sinus, small to medium teeth compared to their width at the base with straight or convex sides, no anthocyanin coloration of veins, a twisted, finely blistered leaf blade, and on the lower side of the leaves, no or a very low density of erect and prostrate hairs,

- the long and often fasciated bunches,
- the ellipsoid berries that are yellow pinkish when fully ripe.

# **Genetic profile**

MicrosatelliteVVS2		VVMD5	VVMD7	VVMD27	VRZAG62	VRZAG79	VVMD25	VVMD28	VVMD32
Allele 1	143	225	249	176	188	256	240	257	255
Allele 2	149	232	253	191	204	258	248	267	271

#### Cultivation and agronomic skills

Patricia INTA has a semi-erect bearing. Under favorable circumstances, this variety is very productive and produce high yields. It is sensitive to magnesium deficiency.

#### **Clonal selection in France**

The only certified Patricia INTA clone carries the number 1364.

#### Phenology

Bud burst: 9 days after Chasselas. Grape maturity: late-season, 5 weeks after Chasselas.


#### Debourrement

#### **Technological potential**


Patricia INTA's bunches are large to very large, long and moderately compact. The berries are medium in size, seedless, with a slightly muscat flavor and a thin skin. The berries turn pink only when they are fully ripe. When the seeds are present, they are soft and not very noticeable.

## Susceptibility to Diseases and Pests

Patricia INTA is very sensitive downy mildew.



Maturité



#### **Bibliographic references**

- Documentary collections of the Centre de Ressources Biologiques de la Vigne de Vassal-Montpellier, INRAE - Institut Agro Montpellier, Marseillan, France.



Plantgrape, all rights reserved, plantgrape.fr, UMT Géno-Vigne® INRAE - IFV - L'Institut Agro Montpellier