


Riparia Gloire de Montpellier



Genetic origin

This is a *Vitis riparia* Michaux selection.

Breeder/breeder and year obtained

L. Vialla and R. Michel, 1880.

Estimated surface area of the French vineyard grafted with this rootstock and main regions of use

17 000 ha . Aquitaine, Val de Loire Midi-Pyrénées, Rhône-Alpes, Bourgogne Franche-Comté, Charentes, Languedoc-Roussillon.

Elements of ampelographic description

The identification is based on:

- the tip of the young shoot that is completely closed, in the shape of a crook with a very high density of erect hairs but no prostrate hairs,
- the green young leaves, with well developed stipules at the base of the petioles,
- the shoots with a horizontal bearing, an even surface, no bloom, an elliptic section, no erect and prostrate hairs, not very noticeable nodes with long, colored on the dorsal side internodes,
- the large, cordate, entire adult leaves, with an widely open U-shaped petiole sinus, an undulate leaf blade between the veins, long teeth compared to their width, with straight sides or with one side slightly convex and one side slightly concave (the three terminal teeth of the central main vein and the two main adjacent lateral veins are longer and more developed), and on the lower side of the leaves, no prostrate hairs and a high density of erect hairs on the veins, more particularly on the bifurcation of the veins,
- the male flowers,
- the long, brown woody shoots with no erect and prostrate hairs.

Evolution of mother vine surfaces

Year	1945	1955	1965	1975	1985	1995	2005	2015
ha	129	216	116	52	27	52	45	30

Genetic profile

Microsatellite	VVS2	VVMD5	VVMD7	VVMD27	VRZAG62	VRZAG79	VVMD25	VVMD28	VVMD32
Allele 1	139	263	251	236	192	256	236	214	236
Allele 2	143	263	264	238	200	260	238	243	236

Resistance to soil pests

Riparia Gloire de Montpellier is very highly tolerant to the root form of phylloxera. It is also quite tolerant to *Meloidogyne hapla* nematodes, but its resistance to *Meloidogyne incognita* and *Meloidogyne arenaria* nematodes is only moderate. It is quite tolerant to *Agrobacterium vitis*.

Aptitudes for vegetative multiplication

Riparia Gloire de Montpellier wood production is good (40 000 à 80 000 m/ha). The internodes are long with an average diameter. The growth of lateral shoot buds is limited and the canes are not very ramified. This rootstock has good cuttings and grafting capacities.

Clonal selection in France

In France, the 5 certified Riparia Gloire de Montpellier clones carry the numbers 1, 142, 186, 1030 and 1162. Among those, the clones multiplied are:

- clone No. 1: 12 ha 24 ares of mother vines producing certified material, in 2017,
- clone No. 142: 3 ha 40 ares of mother vines producing certified material, in 2017,
- clone No. 186: 50 ares of mother vines producing certified material, in 2017,
- clone No. 1030: 13 ha 29 ares of mother vines producing certified material, in 2017.

Datas are extracted from: Les chiffres de la pépinière viticole, 2017, Datas and assesment of FranceAgriMer, may 2018.

Bibliographic references

- Catalogue des variétés et clones de vigne cultivés en France. Collectif, 2007, Ed. IFV, Le Grau-du-Roi, France.
- Documentary collections of the Centre de Ressources Biologiques de la Vigne de Vassal-Montpellier, INRAE - Montpellier SupAgro, Marseillan, France.
- Cépages et vignobles de France, tome 1. P. Galet, 1988, Ed. Dehan, Montpellier, France.

Adaptation to the environment

This rootstock is adapted to acidic or decalcified soils. On the other hand, it is very sensitive to chlorosis and only resists to less than 15% of "total" limestone, 6% of "active" limestone and an IPC of 5. It is well adapted to humidity but is very sensitive to drought and is poorly adapted to compact soils with too much clay. This rootstock absorbs easily potassium, but poorly absorbs magnesium. It may thus promote magnesium deficiency symptoms, dessication of the stems and less acidic musts. This rootstock should be used in gravelly and fairly fertile soils with a sufficient water input. It should not be avoided in poor, dry or superficial soils.

Interaction with the graft and production objectives

The radial trunk growth of this rootstock remains limited. It is said that this rootstock has a "thin leg" which may create substantial diameter differences with the grafts and the requirement for trellising. The development speed of the young plants is good. Riparia Gloire de Montpellier confers a very low vigor and the vine development is reduced. Riparia Gloire de Montpellier promotes floral initiation but given its very weak vigor, the yields obtained are usually limited. This rootstock has an influence on the earliness of the vegetative growth and under adapted conditions, with a sufficient plant density it induces the production of quality fruits. Riparia Gloire de Montpellier has a very good affinity and works well with Cabernet-Sauvignon, Chardonnay, Chenin, Cot, Gros Manseng, Merlot, Sauvignon and Tannat.



*Plantgrape, all rights reserved,
plantgrape.fr, UMT Géno-Vigne®
INRAE - IFV - L'Institut Agro Montpellier*